


# Kinetic Theory of Gases

## Physics 1425 Lecture 31

# Bernoulli's Picture


- Daniel Bernoulli, in 1738, was the first to understand air pressure in terms of molecules—he visualized them shooting around very rapidly in a closed container, supporting a weight as shown by constantly bouncing off the underside of the piston.
- Given more room, they would rush in to fill the new space, just as a gas is observed to do.
- No-one believed him.



[Applet here.](#)

# One Dimensional, One Molecule Gas


- The **molecule** roundtrips in time  $2L/v$ , so it bounces off the piston  $v/2L$  times per sec, each time delivering momentum  $2mv$ , so the **piston** will pick up momentum from this “gas” at rate  $2mv \times v/2L$  per second.
- Force from gas on piston:  
 $F = \text{rate of change of momentum} = mv^2/L$ . An equal opposite force must be supplied from outside to keep the piston at rest.
- Animation!



1-D gas: molecule bounces between ends of cylinder.

# Molecule in a Two-Dimensional Box

- Assume perfectly elastic collisions with all walls.
- The molecule will follow a **zigzag** path, the time between collisions with the same end, say the end at  $x = L$ , is now  $2L/v_x$ , and the momentum transferred per collision is  $2mv_x$ , so the average force on the end is  $mv_x^2/L$ .
- This will still hold good in *three* dimensions.



# $N$ Molecules in an $L \times L \times L$ Cube



- Assume first that we have a very large number  $N$  of molecules bouncing around, so small that they don't hit each other, each follows its own zigzag path.
- The force on the right-hand wall at  $x = L$  is just the sum of the forces from each one, so
  - $F = mv_{x1}^2/L + mv_{x2}^2/L + mv_{x3}^2/L + \dots + mv_{xN}^2/L .$

# Bar Means Average

- The force on the wall depends on the sum

$$v_{x1}^2 + v_{x2}^2 + v_{x3}^2 + \dots + v_{xN}^2$$

- We don't need the *individual* values  $v_{x1}^2$ , etc., just the *average*, written **with a bar**:

$$\overline{v_x^2} = \frac{v_{x1}^2 + v_{x2}^2 + v_{x3}^2 + \dots + v_{xN}^2}{N}$$

- So the force on the wall is:

$$F = \frac{Nm\overline{v_x^2}}{L}$$

# Gas Molecules Have *Random* Velocities...

- Gravity has a negligible effect on the speeds, so **all directions look the same** to the gas molecules.

- This means:  $\overline{v_x^2} = \overline{v_y^2} = \overline{v_z^2}$   
and since the molecular speed squared

$$v^2 = v_x^2 + v_y^2 + v_z^2$$

it follows that

$$\overline{v_x^2} = \frac{1}{3} \overline{v^2}$$

# Relating Pressure to Molecular Energy

- Our gas of molecules is in an  $L \times L \times L$  cube, the force on one wall (which has **area**  $L^2$ ) is

$$F = \frac{Nm\overline{v_x^2}}{L} = \frac{Nm\overline{v^2}}{3L}$$

- So the **pressure**  $P$ , force per unit area, is ( $V = L^3$ )

$$P = \frac{F}{L^2} = \frac{Nm\overline{v^2}}{3L^3} = \frac{2N\left(\frac{1}{2}\overline{mv^2}\right)}{3V}$$

- Now we'll multiply both sides by  $V$  ...



# Finding the Ideal Gas Law

- We've established that  $PV = \frac{2}{3} N \left( \overline{\frac{1}{2} m v^2} \right)$  and we know that for very weakly interacting gases,  $PV = nRT$ . *These two equations must be the same!*
- The equivalence is most simply expressed using Boltzmann's constant,  $k = R/N_A$  ( $= 1.38 \times 10^{-23}$  J/K).

$$PV = nRT = nN_A kT = NkT, \text{ so}$$

$$\overline{\frac{1}{2} m v^2} = \frac{3}{2} kT$$

- Absolute temperature is proportional to average molecular kinetic energy.

# Average Speed of Air Molecules

- Maxwell and co were very smart guys—they figured out accurately the average speed of air molecules *before they had any idea how big the molecules were!*
- They just used

$$PV = \frac{2}{3} N \left( \overline{\frac{1}{2} m v^2} \right) = \frac{1}{3} N m \overline{v^2} = \frac{1}{3} M \overline{v^2}$$

where  $M$  is the total mass of the gas in the box.

# Average Speed of Air Molecules

- Let's see what  $PV = \frac{1}{3}M\overline{v^2}$  gives for the speed of air molecules in this room (we are, of course, averaging here over oxygen and nitrogen plus a tiny amount of other stuff).
- Let's take a **one meter cube**: it will contain about 1.3 kg of air. The pressure  $P = 10^5 \text{ N/m}^2$ , close enough, so

$$PV = 10^5 = \frac{1}{3}(1.3)\overline{v^2}$$

giving the root mean square value  $\sqrt{\overline{v^2}} = 480 \text{ m/s}$ .

# Clicker Question

- Since oxygen, nitrogen and helium all satisfy the same gas law  $PV = nRT$  at room temperature, we conclude that:
  - A. All have the same (rms) root mean square average molecular speed
  - B. All have the same average molecular kinetic energy
  - C. Neither of the above is true.

# Clicker Question

- If we take the average speed of oxygen molecules in this room to be 480 m/s, what would be the average speed of helium atoms that leaked from a balloon into the room?
- A. 480 m/s
- B. 960 m/s
- C. 1360 m/s
- D. 1920 m/s

# The Speed Distribution

- Although the molecules fly freely almost all of the time, they *do* collide occasionally.
- Assuming random elastic collisions, there will be transfer of energy, typically of order  $kT$ , from one to another in a collision.
- The chances of a particular molecule picking up  $kT$   $n$  times in a row is similar to the chances of a coin toss coming up heads  $n$  times in a row.... that is, high energies are *exponentially unlikely*.

# Maxwell's Speed Distribution

- Maxwell did the math precisely, and found the probability of a molecule having a high energy at a given moment did drop exponentially with energy:

$$\text{Probability (speed} = v) \propto e^{-\frac{1}{2}mv^2/kT}$$

meaning that for each extra  $kT$  of energy, the probability of finding a particle with that energy drops by  $1/e \approx 0.37$  —more than a factor of 2, because the average amount picked up per collision is less than  $kT$ .

- (Maxwell's exact result is  $f(v) = 4\pi N \left( \frac{m}{2\pi kT} \right)^{\frac{3}{2}} v^2 e^{-\frac{1}{2}mv^2/kT}$ .)

# Escaping from a Planet

- Maxwell's speed distribution makes it quite easy to predict which gases can escape from planetary atmospheres.
- For the Earth,  $v_{\text{escape}} = 11 \text{ km/sec}$ .
- The upper atmosphere has parts as hot as 1000K.
- The speed distribution includes  $e^{-mv^2/2kT} = e^{-3v_{\text{escape}}^2/2v^2}$  where we've used  $\frac{1}{2}mv^2 = \frac{3}{2}kT$ .
- For  $\text{H}_2$ , at 1000K, the fraction of molecules at escape velocity is of order  $10^{-6}$ , for He  $10^{-12}$ , for  $\text{O}_2$   $10^{-79}$ . This means the  $\text{H}_2$  will escape almost instantly, the He pretty quickly, and the  $\text{O}_2$  never.




# Clicker Question

- Which of the following gases would you expect to be dominant in the Martian atmosphere?
- A.  $\text{H}_2$
  - B. He
  - C.  $\text{N}_2$
  - D.  $\text{H}_2\text{O}$
  - E.  $\text{CO}_2$

# Real Gases

- (This is of course chemistry.)
- **Water** has the three phases as shown on this pressure/temperature graph.
- They meet at the “**triple point**” a definite  $P$  and  $T$ , useful as a reference point in fixing temperatures.
- Freezing and boiling points vary with pressure.
- At the **critical point**, liquid and vapor become the same.




The solid-liquid dividing line slopes *forwards* for almost all substances—water is an exception.

# Vapor Pressure and Humidity

- The  $\text{H}_2\text{O}$  molecules in liquid water strongly attract each other, holding the liquid together. But these molecules are still jiggling around, with a Maxwell speed distribution. This means a fraction of them near the surface are moving fast enough to escape, forming a vapor above the surface.
- In a closed container, with enough water present, an **equilibrium** situation is reached between escaping and returning molecules.

# Water and Vapor in Equilibrium

- In equilibrium in a closed container, the molecules in the vapor have the same average kinetic energy as the air molecules, so exert pressure on the walls of the container proportionate to their numbers.
- This is the **saturated vapor pressure**. It varies with temperature like  $e^{-a/T}$ , not surprising since its origin is molecules fast enough to escape.



# Water Vapor Pressure

- At room temperature, saturated vapor pressure is about 2.5% of atmospheric pressure.
- At 100°C, it **equals** atmospheric pressure: this means small bubbles formed in the liquid by fast moving molecules coming together are no longer crushed by the surrounding atmospheric pressure, the water boils.
- In mountain resorts like Aspen, water boils at a lower temperature, producing inferior tea.

# Relative Humidity and Dew Point

- If water is constantly boiled off in a closed room, it is not possible to establish an equilibrium situation with vapor pressure above the saturated value for that temperature—further water will condense out on the walls, etc. At this point, relative humidity = 100%.
- **Relative humidity =**  
**vapor pressure/saturated vapor pressure**

**Dew point:** temperature at which dew forms—that is, water condenses out as the air cools.